



THE UNIVERSITY *of* EDINBURGH

Edinburgh Research Explorer

RNA-Associated Early-Stage Antiviral Factor Is a Major Component of Lv2 Restriction

Citation for published version:

Marno, K, O'Sullivan, E, Jones, C, Diaz-Delfin, J, Pardieu, C, Sloan, RD & McKnight, Á 2017, 'RNA-Associated Early-Stage Antiviral Factor Is a Major Component of Lv2 Restriction', *Journal of Virology*, vol. 91, no. 10. <https://doi.org/10.1128/JVI.01228-16>

Digital Object Identifier (DOI):

[10.1128/JVI.01228-16](https://doi.org/10.1128/JVI.01228-16)

Link:

[Link to publication record in Edinburgh Research Explorer](#)

Document Version:

Peer reviewed version

Published In:

Journal of Virology

General rights

Copyright for the publications made accessible via the Edinburgh Research Explorer is retained by the author(s) and / or other copyright owners and it is a condition of accessing these publications that users recognise and abide by the legal requirements associated with these rights.

Take down policy

The University of Edinburgh has made every reasonable effort to ensure that Edinburgh Research Explorer content complies with UK legislation. If you believe that the public display of this file breaches copyright please contact openaccess@ed.ac.uk providing details, and we will remove access to the work immediately and investigate your claim.



RNA-associated Early-stage Anti-viral Factor (REAF) is a major component of LV2 restriction

Kelly M. Marno, Eithne O’Sullivan, Christopher E. Jones, Julieta Díaz-Delfín, Claire Pardieu,
Richard D. Sloan and Áine McKnight#

Centre for Immunobiology, Blizard Institute, Barts and The London School of Medicine and
Dentistry, Queen Mary University of London, London, United Kingdom

Running Head: REAF is Lv2

#Address correspondence to Áine McKnight, a.mcknight@qmul.ac.uk

Kelly M. Marno, kelly.marno@gmail.com

Eithne O’Sullivan, eithneosc@gmail.com

Christopher E. Jones, c.e.jones@qmul.ac.uk

Julieta Díaz-Delfín, diazdelfin_j@yahoo.com

Claire Pardieu, c.pardieu@qmul.ac.uk

Richard D. Sloan, r.d.sloan@qmul.ac.uk

Word count (abstract): 189+102

Word count (text): 3241

Abstract

HIV and SIV replication in human cells is restricted at early post-entry steps by host inhibitory factors. We previously described and characterised an early phase restriction of HIV-1 and 2 replication in human cell lines, primary macrophages and PBMCs. The restriction was termed Lentiviral restriction 2 (Lv2). The viral determinants of Lv2 susceptibility mapped to the HIV-2 Env and CA. We subsequently reported a whole genome siRNA screen for factors involved in to HIV which identified RNA-associated Early-stage Anti-viral Factor (REAF). Using HIV-2 chimeras of susceptible and non-susceptible viruses we show here that REAF is a major component of the previously described Lv2. Further studies of the viral CA demonstrate that the CA mutation I73V (previously called I207V), a potent determinant for HIV-2, is a weak determinant of susceptibility for HIV-1. More potent CA determinants for HIV-1 REAF restriction were identified at P38A, N74D, G89V and G94D. These results firmly establish that in HIV-1 CA is a strong determinant of susceptibility to LV2/REAF. Similar to HIV-2 the HIV-1 Env can rescue sensitive CAs from restriction. We conclude that REAF is a major component of the previously described Lv2 restriction.

Importance

Measures taken by the host cell to combat infection drive the evolution of pathogens to counteract or side step them. The study of such virus-host conflicts can point to possible weaknesses in the arsenal of viruses and may lead to the rational design of anti-viral agents. Here we describe our discovery that the host restriction factor REAF fulfils the same criteria previously used to describe Lentiviral restriction (LV2). We show that, like HIV-1 CA, the CA of HIV-1 is a strong determinant of LV2/REAF susceptibility. We illustrate how HIV counteracts LV2/REAF by using an envelope with alternative routes of entry into cells.

Introduction

Infection of cells by human and simian immunodeficiency virus (HIV and SIV) is initiated by binding of the viral envelope (Env) to CD4. Conformational changes in the viral Env expose a site that can interact with a chemokine receptor, either CXCR4 or CCR5, expressed at the cell surface of CD4⁺ T cells and primary macrophages (1, 2). Viruses in general can enter cells through different routes, either directly at the plasma membrane (PM) or through one of a number of endocytic pathways (3). Influenza is a prototypical virus that enters cells through an endocytic route and requires the acid environment of the late endosome to trigger its fusion and entry into cells. Since the mechanism of HIV fusion is pH independent (4) it has been widely assumed that HIV fuses at the PM (5-7). pH independent endocytic entry has recently been

observed (8-15) and is thus a possible mechanism of HIV entry; this however remains a topic of considerable controversy (16, 17). Regardless of the route, once HIV fuses at the plasma membrane the conical core is released into the cytoplasm. The viral genomic RNA is reverse transcribed by the virally encoded RNA/DNA dependent reverse transcriptase (RT), resulting in virally encoded RNA:DNA, single and double stranded (ss and ds)DNA intermediates. The RNase H activity of RT degrades the RNA from these hybrids resulting in ssDNA from which the second DNA strand is synthesised (18, 19). Once reverse transcription is complete the double stranded proviral DNA is processed for integration into the host cell genome.

HIV must overcome many cellular barriers to its replication as it journey to the nucleus to integrate into the host genome (20, 21). Interferon-induced transmembrane proteins (IFITMs) can inhibit virus-cell membrane fusion (22) and the process of reverse transcription itself is also vulnerable. Immediately after initiation, members of the Apolipoprotein B mRNA-editing, enzyme-catalytic, polypeptide-like (APOBEC) family of restriction factors induce deoxycytidine to deoxyuridine mutations in the nascent DNA (23). Further disabling reverse transcription SAMHD1 depletes the dNTP substrates required (24). RNA-associated Early-stage Anti-viral Factor (REAF) was described to inhibit HIV and SIV replication during reverse transcription (25). REAF is intrinsically expressed and provides an initial line of defence against HIV and SIV infection. It associates with reverse transcripts; either ssDNA or RNA:DNA hybrids, however the precise mechanism of its action is not yet understood. A more recently described restriction factor MX2/MXB inhibits replication at a later stage, suppressing nuclear import and proviral formation (26-28). Integration is inhibited by the TRIM28 (KAP1)/SETDB1 complex (29). Once

the provirus is integrated the late phase of the replication cycle begins with the production of viral proteins (30). A plasma membrane located restriction factor tetherin/BST2/CD317, prevents viruses from leaving the cell at the late budding stage of the life cycle (31).

The first Lentiviral restriction factor 1 (Lv1) effective against HIV-1 was identified as rhesus TRIM5 α (32, 33). Lv1/TRIM5 α is species specific and active against HIV-1 in non-human primate cells. TRIM5 α forms a lattice around the capsid (CA) resulting in premature disassembly of the conical core (34). It is not known if Lv2 is species specific. It inhibits HIV-1 and 2 during reverse transcription and susceptibility is determined by the viral CA (35). Lv2 differs from Lv1 in that the Env is an additional determinant of restriction. Approximately half of HIV-1 and HIV-2 viruses are susceptible to Lv2 (35). Lv3 is a post entry restriction to infection of simian MAGI cells by HIV-1 and similar to Lv2 is dependent on fusion events at the cell membrane (36). Lv4 restricts nuclear entry of SIV isolates in human cells (37). A recently described restriction to HIV-1 induced by TLR 7/8 agonist in human monocytes is termed Lv5 (38). So far the identities of Lv3, 4 and 5 are unknown. Here we describe the identification of REAF as a potent component of Lv2.

To identify components of Lv2 we designed a whole genome siRNA screen (20). HeLa-CD4 cells were transfected with an siRNA library targeting 19,121 human genes and then challenged with an HIV-1^{89.6} (MCR) pseudovirus (20). One factor identified was RPRD2 (here called RNA-associated Early-stage Anti-viral Factor; REAF) which we now show fulfils the characteristics used to define Lv2 (35,38).

Materials and Methods

Cells. Buffy coats from seronegative donors were obtained from the National Blood Service (Brentwood, UK). Donors were anonymous and patient consent was not required. Peripheral blood mononuclear cells (PBMC) were prepared by density-gradient centrifugation (Lymphoprep, Axis-Shield). Monocyte-derived macrophages (MDM) were isolated from PBMC using CD14⁺ MACS Microbeads (Miltenyi Biotec) and left to differentiate for 5 days in RPMI 1640/10% foetal calf serum (FCS) and 15ng/ml granulocyte macrophage colony stimulating factor (GM-CSF; Peprotech). HEK 293T, HeLa-CD4, U87-CD4-CXCR4, HeLa EKV and HeLa EKVΔCPSF6-358 cells and their optimal culture conditions have been described previously (39-42).

Preparation of REAF knockdown cells. The pSUPER RNAi system (pSUPER.retro.puro; Oligoengine) was used for expression of shRNA in mammalian cells (43, 44). For REAF knockdown pSUPER.retro.puro(shREAF) was generated by digestion with BglII and HindIII, annealing of the specific primers and ligation. The shRNA target sequences are shown in upper case within the primers listed below:

shREAF-BglII: 5' gatccccCACGTAAGCCCTCAGATGAttcaagagaTCATCTGAGGGCTTACGTGttttta 3'

shREAF-HindIII: 5' agcttaaaaaCACGTAAGCCCTCAGATGAtctcttgaaTCATCTGAGGGCTTACGTGggg 3'

The vector was either transfected directly into HeLa-CD4 cells for transient knockdown or used to generate stable knockdown cell lines. Briefly, retroviruses were produced by co-transfecting

pSUPER.retro.puro(shREAF) with an HIV-1 *gag-pol* expression vector (p8.91) (45) and pMDG VSV-G Env into HEK 293T cells. Supernatant containing virus was harvested after 48 hours and was used to transduce HeLa-CD4 cells under puromycin selection. REAF silencing in transient and stably knocked down cells was confirmed by Western blot.

Plasmids and virus production. The infectious molecular clone for HIV-1^{89.6} was obtained from the Centre for AIDS Research (NIBSC, UK). Infectious full-length and chimeric HIV clones were prepared by polyethylenimine (PEI; Polysciences) or Lipofectamine 2000 (Invitrogen) transfection of HEK 293T cells. The virus named in parentheses for each pseudotype denotes the Env used.

Production of CA mutant viruses. HIV-1 CA mutants were generated by site directed mutagenesis (SDM) of the HIV-1^{NL4.3}-derived viral clone pBR-NL43-IRES-eGFP (46) with further modification to introduce stop codons in the first and third codons of the Env coding sequence. HIV-1 pseudovirus particles were produced by PEI transfection of HEK 293T cells using a 1:1 molar ratio of viral plasmid to MCR/MCN/VSV-G/NL4.3 Env expression plasmid.

Western blot. SDS-PAGE separated proteins were detected with the primary rabbit polyclonal antibody against REAF (Eurogentec) or GAPDH (loading control; Abcam) followed by horseradish peroxidase-conjugated donkey α -rabbit antibody (GE Healthcare). Protein was visualised using a chemiluminescence kit (ECL; GE Healthcare).

siRNA transfection and infection with replication competent virus. HeLa-CD4 cells were seeded at 2.5×10^4 cells/well in 24-well plates. siRNA transfection (30nM) was performed using HiPerfect (QIAGEN) according to the manufacturer's instructions using the following sequences:

siREAF: 5' CACGTAAGCCCTCAGATGATA 3'

siCB: 5' ACAGCAAATTCCATCGTGT 3'

72 hours after siRNA transfection, cells were challenged with virus for up to 5 hr. Infection was assessed up to 48 hours by intracellular p24 staining.

***In situ* immunostaining for p24 antigen.** Infected cells were fixed with cold (-20°C) methanol:acetone (1:1), washed with PBS then immunostained for p24 using mouse anti-HIV-1 p24 monoclonal antibodies EVA365 and 366 (NIBSC, UK) (1:50) as previously described (47). Infected cell foci stained blue (regarded as foci of infection (FFU/ml)) and were quantitated by light microscopy.

Statistical analysis. The results presented are derived from a minimum of three independent experiments performed in duplicate at minimum. Differences between two treatments were tested for statistical significance using unpaired two-tailed *t*-Tests. * denotes $p < 0.05$, *n.s.* not significant.

Results

The HIV-2 molecular determinants of Lv2 restriction were previously mapped using two HIV-2 molecular clones of viruses derived from the same patient, HIV-2^{MCR} and HIV-2^{MCN}, which are differentially sensitive to Lv2. A gene swapping approach between the viruses identified the *gag* and *env* genes as critical determinants of Lv2 restriction (39). These chimeric viruses (shown schematically in Fig. 1A) were tested to determine if they had the same pattern of susceptibility to REAF.

HeLa-CD4 cells were knocked down for REAF using specific (siREAF) or non-targeting control siRNA (cyclophilin B, siCB) (Fig. 1B). Fig. 1C shows viral rescue in HeLa-CD4 cells following treatment with siREAF and compared with cells treated with siCB. Repeat experiments consistently show, as expected for a virus highly sensitive to Lv2 (39), that the HIV-2^{MCR} virus is potently rescued in comparison to HIV-2^{MCN} (50 fold vs 10 fold; $p = 0.004$). When the *env* and *gag* from the restricted HIV-2^{MCR} was inserted in place of the relatively insensitive HIV-2^{MCN} *env* and *gag* (HIV-2^{MCN_{mcr} env+gag}), greater sensitivity of this virus to REAF was observed (66 fold). In the reciprocal experiment, where the *env* and *gag* from HIV-2^{MCN} replaced the HIV-2^{MCR} genes (HIV-2^{MCR_{mcr} env+gag}), the resulting chimera was only rescued 3 fold ($p < 0.001$). These results for susceptibility to REAF are consistent with the Lv2 phenotype previously described (39). A single point mutation in HIV-2^{MCR} CA at position 73 is known to be a critical determinant of Lv2 restriction (previously labelled position 207). Fig. 1C shows that HIV-2^{MCR CA 173V} is rescued only 12 fold from REAF restriction compared to 50 fold for wild type HIV-2^{MCR} ($p = 0.003$).

To confirm these results and for further experiments we generated HeLa-CD4 cell lines permanently expressing shRNA specific for REAF mRNA. Western blot (WB) analysis shows that the HeLa-CD4-shREAF cells expressed much less REAF protein than the parental HeLa-CD4 cells

(Fig. 1D). The phenotype of knockdown of REAF in this cell line was confirmed using HIV-2^{MCR} and HIV-2^{MCR CA I73V}. The HIV-2^{MCR} virus was restricted 326 fold compared to 33 fold for the HIV-2^{MCR CA I73V} (Fig. 1E; $p = 0.016$).

We previously reported that Lv2 was active in HeLa-CD4, human primary PBMC and MDM but not in U87-CD4-CXCR4 cells (39). REAF mRNA (data not shown) and protein is present in MDM and to much lower levels in PBMC (Fig. 1F) while WB analysis shows it to be barely detectable in U87-CD4-CXCR4 (Fig. 1G).

As previously reported for Lv2 (39) and further demonstrated here, the HIV-2 determinants for REAF are Env and CA (specifically amino acid 73). We sought to identify the determinants of REAF restriction for HIV-1. Fig. 2A shows that, compared to HIV-1^{89.6}, HIV-1^{NL4.3} is more resistant to REAF restriction (3 fold vs 21 fold; $p < 0.001$). We used HIV-1^{NL4.3} to further establish if the equivalent HIV-2 CA mutation 73 plays a role in Lv2/REAF restriction in HIV-1. Using SDM we generated HIV-1^{NL4.3 CA I73V}. Both wild type and mutant CA were pseudotyped with HIV-2^{MCR} Env and/or HIV-1^{NL4.3} Env and tested for their susceptibility to REAF using the HeLa-CD4-shREAF cell line. Fig. 2B shows that wild type HIV-1^{NL4.3} CA is only weakly susceptible to REAF when pseudotyped with HIV-1^{NL4.3} Env. However when the CA is mutated (HIV-1^{NL4.3 CA I73V} (NL4.3)) the restriction is more potent but still relatively weak compared to HIV-2 (15 fold, compare to Fig. 1E for HIV-2). The CA I73V was further restricted when pseudotyped with an HIV-2^{MCR} Env (21 fold, $p = 0.03$), but not with HIV-2^{MCN} Env (16 fold, $p = n.s.$). Thus CA amino acid 73 and Env are determinants of Lv2/REAF restriction for both HIV-1 and 2.

222 The CA amino acid at position 73 lies in the binding domain of the cleavage and polyadenylation
223 specific factor 6 (CPSF6) protein (48). This is of particular interest as the adjacent CA mutation
224 (N74D) has been shown to affect the sensitivity of HIV-1 to depletion of RanBP2, Nup153 and
225 TNPO3 nuclear pore proteins (49, 50).

226 N74D is an HIV-1 escape mutant that was generated by passage of HIV-1^{NL4.3} in cells expressing
227 an artificially mutated CPSF6-358 that perturbs HIV-1 nuclear entry (50). CPSF6 is a pre-mRNA
228 processing protein that shuttles between the nucleus and the cytoplasm (51). The mutant form
229 CPSF6-358 lacks a C-terminal nuclear-targeting arginine/serine (RS)-rich domain and so is
230 confined to the cytoplasm and restricts HIV-1 before nuclear entry (50, 52, 53).

231 We tested whether the Lv2/REAF HIV-2 CA determinant I73V was similar to N74D with respect
232 to resistance to CPSF6-358. Another CA mutation, P38A, which was mutated outside the CPSF6
233 CA binding region was included as a negative control (54). HeLa-CD4 target cells permanently
234 expressing the mutant CPSF6-358 (HeLa EKVΔCPSF6-358) or vector alone (HeLa EKV) (42, 48)
235 were challenged with pseudotypes carrying a mutant or wild type CA. Fig. 2C shows that, as
236 expected, infection of the pseudotypes with wild type and P38A CA are inhibited (62 and 73
237 fold respectively). However both CA mutants, HIV-1^{NL4.3 CA I73V} (VSV-G) and HIV-1^{NL4.3 CA N74D} (VSV-
238 G), were resistant to CPSF6-358 (3.3 and 1.6 fold; both $p < 0.001$).

239

240 Given this similarity in resistance to CPSF6-358 we also sought to determine whether, similar to
241 I73V, N74D is more susceptible than wild type virus to REAF restriction. HeLa-CD4 and HeLa-
242 CD4-shREAF cells were challenged by pseudotypes with either wild type or mutated CA. Fig. 3A
243 shows that HIV-1^{NL4.3} (MCR) is rescued 13 fold. Surprisingly HIV-1^{NL4.3 CA N74D} (MCR) is even more

restricted and is rescued 53 fold ($p < 0.001$), suggesting CA N74D is a more potent determinant of REAF restriction in HIV-1

As well as being less sensitive to CPSF6-358, the CA N74D mutation has a more stable conical core which results in delayed disassembly and reverse transcription (55). It is thought that optimal disassembly of the conical core is required for successful infection as mutations interfering with core stability often result in a disturbance of reverse transcription kinetics (54, 56-58). We previously reported that REAF was transiently down modulated shortly after infection (25). We hypothesised that unstable capsids will prematurely disassemble and expose reverse transcripts to REAF. The corollary of this is that capsids that disassemble too late will miss the window of time where REAF is absent. To test this hypothesis that capsid stability was a determinant of REAF/Lv2 restriction we investigated the REAF susceptibility of CA mutations with varying CA stability. P38A, in contrast to N74D, is highly unstable (54, 59) while G94D is unaffected (55). The G89V mutation in the cyclophilin binding loop was chosen because it has previously been shown to affect sensitivity to host restriction factors (42, 54, 57, 58). P38A is also distinct from N74D in that it is sensitive to CPSF6-358 (Fig. 2C). Fig. 3C, D and E show the infectivity of all three capsid mutants. HIV-1^{NL4.3 CA P38A} (MCR), HIV-1^{NL4.3 CA G89V} (MCR) and HIV-1^{NL4.3 CA G94D} were severely compromised on HeLa-CD4 cells (titre of <500 FFU/ml; compared to wild type CA, Fig. 3B). However their replication was rescued in the absence of REAF similar to HIV-1^{NL4.3 CA N74D} (MCR) (Fig. 3F; all $p < 0.001$).

We showed above that in addition to CA, HIV-1 Env also confers sensitivity to REAF (Fig. 2B). We previously determined that HIV-2^{M^{CN}} Env has the ability to overcome Lv2 (35, 39). If REAF is Lv2

the HIV-2^{MCN} Env would overcome REAF restriction. HeLa-CD4-shREAF cells were challenged with HIV-1^{NL4.3} with a wild type CA pseudotyped with HIV-2^{MCN} Env. As expected the wild type HIV-1^{NL4.3} CA was only slightly restricted and the HIV-2^{MCN} Env could reduce this to a small but significant degree (5.6 to 3.1 fold; $p = 0.007$) (Fig. 4A). In contrast the HIV-1^{NL4.3} strains carrying REAF sensitive CA mutations N74D, P38A, G89V and G94V were potently rescued with the HIV-2^{MCN} Env (Fig. 4B-E, all p -values < 0.001) further confirming that REAF and Lv2 are similarly rescued by viral Env.

Discussion

Here we show that REAF is a major component of the previously described restriction Lv2 (35, 60, 61). HIV chimeric viruses and mutants that delineate susceptible and resistant clones demonstrate that Lv2 and REAF are indistinguishable. Both Lv2 and REAF restriction activity are molecularly determined by the viral Env and CA. The CA amino acid at position 73 in HIV-2 was a crucial determinant of Lv2 and confirmed here for REAF (39). The equivalent amino acid at position 73 also affects HIV-1 susceptibility to REAF. Although the effects of I73V are statistically significant they are much weaker compared to HIV-2. Indeed, the same amino acid substitution rendered HIV-2^{MCN} *less* susceptible but HIV-1^{NL4.3} *more* susceptible to REAF. This led us to the hypothesis that the overall structure or stability of the CA rather than precise molecular interactions is critical. It has been proposed that disassembly too early prior to localisation at the nuclear pore would result in exposure to restriction factors and premature termination of reverse transcription (19, 62-65). Indeed it has been proposed that HIV-1 disassembly involves a regulated collapse of the conical core which protects viral reverse transcription complexes (66).

Also it has been shown that disassembly occurs within an hour of fusion and is facilitated by reverse transcription (67-69). We have observed that cellular REAF is reduced within one hour of viral challenge, but importantly, levels are rapidly replenished an hour later (25). REAF associates with viral nucleic acid and restricts replication during reverse transcription (25). We therefore suggested that the temporary reduction in REAF protein level allows viruses to reverse transcribe in the absence of REAF associated activity. This model is in keeping with the notion that CA stability is a determinant of REAF associated restriction. We further tested the hypothesis that CA stability is a determinant of REAF susceptibility using already well characterised mutants with different CA stabilities. Compared to I73V (20 fold) more potent effects (more than 50 fold) were observed with mutants P38A (unstable), N74D (hyperstable) and G94D (unaffected) (55). Since all these CA mutations result in greater susceptibility to REAF associated activity despite having divergent CA stabilities we cannot conclude that CA stability is a major determinant of susceptibility to REAF. Regardless we show that the CA is a strong determinant of REAF restriction in HIV-1. Given the multifunctional role of CA in HIV-1 replication (70), understanding the role of CA in susceptibility to REAF associated activity may shed light on more specific interactions of CA with host cell factors required for efficient infectivity of HIV.

It is highly controversial whether or not infectious HIV conical cores enter the cytoplasm after fusion at the plasma membrane or through an endocytic route (8-17, 35, 60). Our previous observation of Lv2 suggests that, although either route is possible, infection is more successful when the virus fuses at the plasma membrane and avoids an endocytic entry pathway (35, 60).

Our results here confirm that, like Lv2, the choice of entry route as determined by the viral Env is a determinant of susceptibility to REAF associated activity. HIV-1^{NL4.3} can be rendered sensitive to REAF if pseudotyped with HIV-2^{MCR} while HIV-2^{MCN} Env does not. These previously characterised HIV-2 Envs fuse either at the plasma membrane (MCN) or via an endocytic route (MCR) [35,38]. Furthermore all CA mutants which were rendered highly sensitive to REAF associated activity were protected when pseudotyped with an MCN Env.

We propose that, similar to Lv2, REAF may be more active against viruses attempting to access the cytoplasm via an endosome. Therefore, fusion at the plasma membrane is a more efficient replication pathway. However we cannot eliminate the possibility that some viruses will bypass REAF associated activity regardless of entry route, for example if they have a conical core without the sensitivity conferring mutations described here. Future studies that address the role of Env and CA in determining REAF associated restriction will shed light on these early host cell interactions in the early life cycle of HIV.

Acknowledgements

The monoclonal antibodies to p24 (EVA365 and 366) were provided by the EU Programme EVA Centre for AIDS Reagents, NIBSC, UK (AVIP Contract Number LSHP-CT-2004-503487). We thank Laura Hilditch and Greg Towers for providing HeLa cells expressing CPSF6-358. Thanks to Uta Von Schwedler and Wesley Sundquist for the very kind donation of HIV-1-NL4.3 CA mutants. We would like to remember Uta who is no longer with us to see the publication of this paper.

Author contributions

Designed experiments; ÁMcK, KMM, RDS; Performed experiments; ÁMcK, KM, EOS, CEJ, JDD, CP and RDS; Analysed data; ÁMcK, KMM, RDS. Wrote the manuscript; ÁMcK and KMM.

Funding Information

This work was supported by The Rosetrees Trust (M275 and M275-F1) and a Queen Mary University of London, Life Sciences Institute PhD studentship (C. E. J). The funders had no role in study design, data collection and interpretation, or the decision to submit the work for publication.

References

1. **Clapham PR, McKnight A.** 2002. Cell surface receptors, virus entry and tropism of primate lentiviruses. *J Gen Virol* **83**:1809-1829.
2. **Doms RW, Trono D.** 2000. The plasma membrane as a combat zone in the HIV battlefield. *Genes Dev* **14**:2677-2688.
3. **Marsh M, Helenius A.** 2006. Virus entry: open sesame. *Cell* **124**:729-740.
4. **McClure MO, Sommerfelt MA, Marsh M, Weiss RA.** 1990. The pH independence of mammalian retrovirus infection. *J Gen Virol* **71 (Pt 4)**:767-773.
5. **Maddon PJ, McDougal JS, Clapham PR, Dalglish AG, Jamal S, Weiss RA, Axel R.** 1988. HIV infection does not require endocytosis of its receptor, CD4. *Cell* **54**:865-874.
6. **Pelchen-Matthews A, Clapham P, Marsh M.** 1995. Role of CD4 endocytosis in human immunodeficiency virus infection. *J Virol* **69**:8164-8168.

- 354 7. **Stein BS, Gowda SD, Lifson JD, Penhallow RC, Bensch KG, Engleman EG.** 1987. pH-independent
355 HIV entry into CD4-positive T cells via virus envelope fusion to the plasma membrane. *Cell*
356 **49**:659-668.
- 357 8. **Vidricaire G, Tremblay MJ.** 2007. A clathrin, caveolae, and dynamin-independent endocytic
358 pathway requiring free membrane cholesterol drives HIV-1 internalization and infection in
359 polarized trophoblastic cells. *J Mol Biol* **368**:1267-1283.
- 360 9. **de la Vega M, Marin M, Kondo N, Miyauchi K, Kim Y, Epand RF, Epand RM, Melikyan GB.** 2011.
361 Inhibition of HIV-1 endocytosis allows lipid mixing at the plasma membrane, but not complete
362 fusion. *Retrovirology* **8**:99.
- 363 10. **Carter GC, Bernstone L, Baskaran D, James W.** 2011. HIV-1 infects macrophages by exploiting an
364 endocytic route dependent on dynamin, Rac1 and Pak1. *Virology* **409**:234-250.
- 365 11. **Clotet-Codina I, Bosch B, Senserrich J, Fernández-Figueras MT, Peña R, Ballana E, Bofill M,**
366 **Clotet B, Esté JA.** 2009. HIV endocytosis after dendritic cell to T cell viral transfer leads to
367 productive virus infection. *Antiviral Res* **83**:94-98.
- 368 12. **Daecke J, Fackler OT, Dittmar MT, Kräusslich HG.** 2005. Involvement of clathrin-mediated
369 endocytosis in human immunodeficiency virus type 1 entry. *J Virol* **79**:1581-1594.
- 370 13. **Maréchal V, Prevost MC, Petit C, Perret E, Heard JM, Schwartz O.** 2001. Human
371 immunodeficiency virus type 1 entry into macrophages mediated by macropinocytosis. *J Virol*
372 **75**:11166-11177.
- 373 14. **Sloan RD, Kuhl BD, Mesplède T, Münch J, Donahue DA, Wainberg MA.** 2013. Productive entry
374 of HIV-1 during cell-to-cell transmission via dynamin-dependent endocytosis. *J Virol* **87**:8110-
375 8123.
- 376 15. **Miyauchi K, Kim Y, Latinovic O, Morozov V, Melikyan GB.** 2009. HIV enters cells via endocytosis
377 and dynamin-dependent fusion with endosomes. *Cell* **137**:433-444.

- 378 16. **Herold N, Müller B, Kräusslich HG.** 2015. Reply to "Can HIV-1 entry sites be deduced by
379 comparing bulk endocytosis to functional readouts for viral fusion?" J Virol **89**:2986-2987.
- 380 17. **Marin M, Melikyan GB.** 2015. Can HIV-1 entry sites be deduced by comparing bulk endocytosis
381 to functional readouts for viral fusion? J Virol **89**:2985.
- 382 18. **Peterlin BM, Trono D.** 2003. Hide, shield and strike back: how HIV-infected cells avoid immune
383 eradication. Nat Rev Immunol **3**:97-107.
- 384 19. **Campbell EM, Hope TJ.** 2015. HIV-1 capsid: the multifaceted key player in HIV-1 infection. Nat
385 Rev Microbiol **13**:471-483.
- 386 20. **Liu L, Oliveira NM, Cheney KM, Pade C, Dreja H, Bergin AM, Borgdorff V, Beach DH, Bishop CL,**
387 **Dittmar MT, McKnight A.** 2011. A whole genome screen for HIV restriction factors.
388 Retrovirology **8**:94.
- 389 21. **Simon V, Bloch N, Landau NR.** 2015. Intrinsic host restrictions to HIV-1 and mechanisms of viral
390 escape. Nat Immunol **16**:546-553.
- 391 22. **Lu J, Pan Q, Rong L, He W, Liu SL, Liang C.** 2011. The IFITM proteins inhibit HIV-1 infection. J
392 Virol **85**:2126-2137.
- 393 23. **Bishop KN, Holmes RK, Sheehy AM, Malim MH.** 2004. APOBEC-mediated editing of viral RNA.
394 Science **305**:645.
- 395 24. **Laguet N, Sobhian B, Casartelli N, Ringard M, Chable-Bessia C, Ségéral E, Yatim A, Emiliani**
396 **S, Schwartz O, Benkirane M.** 2011. SAMHD1 is the dendritic- and myeloid-cell-specific HIV-1
397 restriction factor counteracted by Vpx. Nature **474**:654-657.
- 398 25. **Marno KM, Ogunkolade BW, Pade C, Oliveira NM, O'Sullivan E, McKnight Á.** 2014. Novel
399 restriction factor RNA-associated early-stage anti-viral factor (REAF) inhibits human and simian
400 immunodeficiency viruses. Retrovirology **11**:3.

- 401 26. **Goujon C, Moncorgé O, Bauby H, Doyle T, Ward CC, Schaller T, Hué S, Barclay WS, Schulz R,**
402 **Malim MH.** 2013. Human MX2 is an interferon-induced post-entry inhibitor of HIV-1 infection.
403 *Nature* **502**:559-562.
- 404 27. **Kane M, Yadav SS, Bitzegeio J, Kutluay SB, Zang T, Wilson SJ, Schoggins JW, Rice CM,**
405 **Yamashita M, Hatzioannou T, Bieniasz PD.** 2013. MX2 is an interferon-induced inhibitor of HIV-
406 1 infection. *Nature* **502**:563-566.
- 407 28. **Liu Z, Pan Q, Ding S, Qian J, Xu F, Zhou J, Cen S, Guo F, Liang C.** 2013. The interferon-inducible
408 MxB protein inhibits HIV-1 infection. *Cell Host Microbe* **14**:398-410.
- 409 29. **Allouch A, Di Primio C, Alpi E, Lusic M, Arosio D, Giacca M, Cereseto A.** 2011. The TRIM family
410 protein KAP1 inhibits HIV-1 integration. *Cell Host Microbe* **9**:484-495.
- 411 30. **Goff SP.** 2007. Host factors exploited by retroviruses. *Nat Rev Microbiol* **5**:253-263.
- 412 31. **Neil SJ, Zang T, Bieniasz PD.** 2008. Tetherin inhibits retrovirus release and is antagonized by HIV-
413 1 Vpu. *Nature* **451**:425-430.
- 414 32. **Towers G, Bock M, Martin S, Takeuchi Y, Stoye JP, Danos O.** 2000. A conserved mechanism of
415 retrovirus restriction in mammals. *Proc Natl Acad Sci U S A* **97**:12295-12299.
- 416 33. **Cowan S, Hatzioannou T, Cunningham T, Muesing MA, Gottlinger HG, Bieniasz PD.** 2002.
417 Cellular inhibitors with Fv1-like activity restrict human and simian immunodeficiency virus
418 tropism. *Proc Natl Acad Sci U S A* **99**:11914-11919.
- 419 34. **Pertel T, Hausmann S, Morger D, Züger S, Guerra J, Lascano J, Reinhard C, Santoni FA, Uchil PD,**
420 **Chatel L, Bisiaux A, Albert ML, Strambio-De-Castillia C, Mothes W, Pizzato M, Grütter MG,**
421 **Luban J.** 2011. TRIM5 is an innate immune sensor for the retrovirus capsid lattice. *Nature*
422 **472**:361-365.

- 423 35. **Marchant D, Neil SJ, Aubin K, Schmitz C, McKnight A.** 2005. An envelope-determined, pH-
424 independent endocytic route of viral entry determines the susceptibility of human
425 immunodeficiency virus type 1 (HIV-1) and HIV-2 to Lv2 restriction. *J Virol* **79**:9410-9418.
- 426 36. **Pineda MJ, Orton BR, Overbaugh J.** 2007. A TRIM5alpha-independent post-entry restriction to
427 HIV-1 infection of macaque cells that is dependent on the path of entry. *Virology* **363**:310-318.
- 428 37. **Pizzato M, Neagu M, Pertel T, Firrito C, Ziglio S, Zufferey M, Berthoux L, Luban J.** 2013. Lv4, an
429 activity that restricts nuclear entry of SIVMAC/SIVSM in human blood cells, abstr *Frontiers of*
430 *Retrovirology: Complex retroviruses, retroelements and their hosts*, Cambridge,
- 431 38. **Hofmann H, Vanwalscappel B, Bloch N, Landau NR.** 2016. TLR7/8 agonist induces a post-entry
432 SAMHD1-independent block to HIV-1 infection of monocytes. *Retrovirology* **13**:83.
- 433 39. **Schmitz C, Marchant D, Neil SJ, Aubin K, Reuter S, Dittmar MT, McKnight A.** 2004. Lv2, a novel
434 postentry restriction, is mediated by both capsid and envelope. *J Virol* **78**:2006-2016.
- 435 40. **Deng HK, Unutmaz D, KewalRamani VN, Littman DR.** 1997. Expression cloning of new receptors
436 used by simian and human immunodeficiency viruses. *Nature* **388**:296-300.
- 437 41. **Salahuddin SZ, Markham PD, Wong-Staal F, Franchini G, Kalyanaraman VS, Gallo RC.** 1983.
438 Restricted expression of human T-cell leukemia--lymphoma virus (HTLV) in transformed human
439 umbilical cord blood lymphocytes. *Virology* **129**:51-64.
- 440 42. **Rasaiyaah J, Tan CP, Fletcher AJ, Price AJ, Blondeau C, Hilditch L, Jacques DA, Selwood DL,**
441 **James LC, Noursadeghi M, Towers GJ.** 2013. HIV-1 evades innate immune recognition through
442 specific cofactor recruitment. *Nature* **503**:402-405.
- 443 43. **Brummelkamp TR, Bernards R, Agami R.** 2002. A system for stable expression of short
444 interfering RNAs in mammalian cells. *Science* **296**:550-553.
- 445 44. **Brummelkamp TR, Bernards R, Agami R.** 2002. Stable suppression of tumorigenicity by virus-
446 mediated RNA interference. *Cancer Cell* **2**:243-247.

447 45. **Naldini L, Blomer U, Gallay P, Ory D, Mulligan R, Gage FH, Verma IM, Trono D.** 1996. In vivo
448 gene delivery and stable transduction of nondividing cells by a lentiviral vector. *Science* **272**:263-
449 267.

450 46. **Schindler M, Münch J, Kutsch O, Li H, Santiago ML, Bibollet-Ruche F, Müller-Trutwin MC,**
451 **Novembre FJ, Peeters M, Courgnaud V, Bailes E, Roques P, Sodora DL, Silvestri G, Sharp PM,**
452 **Hahn BH, Kirchhoff F.** 2006. Nef-mediated suppression of T cell activation was lost in a lentiviral
453 lineage that gave rise to HIV-1. *Cell* **125**:1055-1067.

454 47. **McKnight A, Clapham PR, Weiss RA.** 1994. HIV-2 and SIV infection of nonprimate cell lines
455 expressing human CD4: restrictions to replication at distinct stages. *Virology* **201**:8-18.

456 48. **Price AJ, Fletcher AJ, Schaller T, Elliott T, Lee K, KewalRamani VN, Chin JW, Towers GJ, James**
457 **LC.** 2012. CPSF6 defines a conserved capsid interface that modulates HIV-1 replication. *PLoS*
458 *Pathog* **8**:e1002896.

459 49. **Schaller T, Ocwieja KE, Rasaiyaah J, Price AJ, Brady TL, Roth SL, Hué S, Fletcher AJ, Lee K,**
460 **KewalRamani VN, Noursadeghi M, Jenner RG, James LC, Bushman FD, Towers GJ.** 2011. HIV-1
461 capsid-cyclophilin interactions determine nuclear import pathway, integration targeting and
462 replication efficiency. *PLoS Pathog* **7**:e1002439.

463 50. **Lee K, Ambrose Z, Martin TD, Oztop I, Mulky A, Julias JG, Vandegraaff N, Baumann JG, Wang R,**
464 **Yuen W, Takemura T, Shelton K, Taniuchi I, Li Y, Sodroski J, Littman DR, Coffin JM, Hughes SH,**
465 **Unutmaz D, Engelman A, KewalRamani VN.** 2010. Flexible use of nuclear import pathways by
466 HIV-1. *Cell Host Microbe* **7**:221-233.

467 51. **Ruepp MD, Aringhieri C, Vivarelli S, Cardinale S, Paro S, Schümperli D, Barabino SM.** 2009.
468 Mammalian pre-mRNA 3' end processing factor CF Im68 functions in mRNA export. *Mol Biol*
469 *Cell* **20**:5211-5223.

- 470 52. **Dettwiler S, Aringhieri C, Cardinale S, Keller W, Barabino SM.** 2004. Distinct sequence motifs
471 within the 68-kDa subunit of cleavage factor Im mediate RNA binding, protein-protein
472 interactions, and subcellular localization. *J Biol Chem* **279**:35788-35797.
- 473 53. **Rüegsegger U, Blank D, Keller W.** 1998. Human pre-mRNA cleavage factor Im is related to
474 spliceosomal SR proteins and can be reconstituted in vitro from recombinant subunits. *Mol Cell*
475 **1**:243-253.
- 476 54. **Forshey BM, von Schwedler U, Sundquist WI, Aiken C.** 2002. Formation of a human
477 immunodeficiency virus type 1 core of optimal stability is crucial for viral replication. *J Virol*
478 **76**:5667-5677.
- 479 55. **Hulme AE, Kelley Z, Okocha EA, Hope TJ.** 2015. Identification of capsid mutations that alter the
480 rate of HIV-1 uncoating in infected cells. *J Virol* **89**:643-651.
- 481 56. **Fitzon T, Leschonsky B, Bieler K, Paulus C, Schröder J, Wolf H, Wagner R.** 2000. Proline residues
482 in the HIV-1 NH2-terminal capsid domain: structure determinants for proper core assembly and
483 subsequent steps of early replication. *Virology* **268**:294-307.
- 484 57. **Li Y, Kar AK, Sodroski J.** 2009. Target cell type-dependent modulation of human
485 immunodeficiency virus type 1 capsid disassembly by cyclophilin A. *J Virol* **83**:10951-10962.
- 486 58. **Shah VB, Shi J, Hout DR, Oztop I, Krishnan L, Ahn J, Shotwell MS, Engelman A, Aiken C.** 2013.
487 The host proteins transportin SR2/TNPO3 and cyclophilin A exert opposing effects on HIV-1
488 uncoating. *J Virol* **87**:422-432.
- 489 59. **von Schwedler UK, Stray KM, Garrus JE, Sundquist WI.** 2003. Functional surfaces of the human
490 immunodeficiency virus type 1 capsid protein. *J Virol* **77**:5439-5450.
- 491 60. **Harrison IP, McKnight A.** 2011. Cellular entry via an actin and clathrin-dependent route is
492 required for Lv2 restriction of HIV-2. *Virology* **415**:47-55.

61. **Reuter S, Kaumanns P, Buschhorn SB, Dittmar MT.** 2005. Role of HIV-2 envelope in Lv2-mediated restriction. *Virology* **332**:347-358.
62. **Lahaye X, Satoh T, Gentili M, Cerboni S, Conrad C, Hurbain I, El Marjou A, Lacabartz C, Lelièvre JD, Manel N.** 2013. The capsids of HIV-1 and HIV-2 determine immune detection of the viral cDNA by the innate sensor cGAS in dendritic cells. *Immunity* **39**:1132-1142.
63. **Landau NR.** 2014. The innate immune response to HIV-1: to sense or not to sense. *DNA Cell Biol* **33**:271-274.
64. **Towers GJ, Noursadeghi M.** 2014. Interactions between HIV-1 and the cell-autonomous innate immune system. *Cell Host Microbe* **16**:10-18.
65. **Gao D, Wu J, Wu YT, Du F, Aroh C, Yan N, Sun L, Chen ZJ.** 2013. Cyclic GMP-AMP synthase is an innate immune sensor of HIV and other retroviruses. *Science* **341**:903-906.
66. **Fassati A, Goff SP.** 2001. Characterization of intracellular reverse transcription complexes of human immunodeficiency virus type 1. *J Virol* **75**:3626-3635.
67. **Hulme AE, Perez O, Hope TJ.** 2011. Complementary assays reveal a relationship between HIV-1 uncoating and reverse transcription. *Proc Natl Acad Sci U S A* **108**:9975-9980.
68. **Kutluay SB, Perez-Caballero D, Bieniasz PD.** 2013. Fates of retroviral core components during unrestricted and TRIM5-restricted infection. *PLoS Pathog* **9**:e1003214.
69. **Yang Y, Fricke T, Diaz-Griffero F.** 2013. Inhibition of reverse transcriptase activity increases stability of the HIV-1 core. *J Virol* **87**:683-687.
70. **Rihn SJ, Wilson SJ, Loman NJ, Alim M, Bakker SE, Bhella D, Gifford RJ, Rixon FJ, Bieniasz PD.** 2013. Extreme genetic fragility of the HIV-1 capsid. *PLoS Pathog* **9**:e1003461.

Figure legends

Figure 1

REAF is an important component of Lv2. A) A schematic representation of the HIV-2 molecular clones, chimeric viruses and site directed mutagenesis (*) used to map the determinants of REAF restriction. B) Western blot of HeLa-CD4 cell lysate following REAF siRNA knockdown compared with non-targeting control (siCB). GAPDH is added as a loading control. C) Titration of constructs on HeLa-CD4 cells transiently transfected with siREAF showing fold change compared to cells transfected with siCB control (compared with HIV-2^{MCR}: HIV-2^{MCN} $p = 0.004$, HIV-2^{MCNmcr env+gag} $p = n.s.$, HIV-2^{MCRmcr env+gag} $p < 0.001$, HIV-2^{MCR CA 173V} $p = 0.003$). D) WB of knockdown of REAF in HeLa-CD4-shREAF cells compared to HeLa-CD4 cells. GAPDH is added as a loading control. E) Fold change for HeLa-CD4-shREAF cells infected with HIV-2^{MCR} and HIV-2^{MCR CA 173V} confirms the Lv2 phenotype in the stable knockdown cells ($p = 0.016$). F) WB of REAF levels in MDM and PBMC compared to HeLa-CD4 cells. GAPDH is added as a loading control. G) WB of REAF levels in U87-CD4-CXCR4 cells compared to HeLa-CD4 cells. GAPDH is added as a loading control.

Figure 2

The HIV-1 CA determinants of REAF associated restriction are the same as those for HIV-2 and are in the CPSF6 binding pocket. A) A comparison of the susceptibility of HIV-1^{NL4.3} and HIV-1^{89.6} following transient knockdown of REAF by siRNA ($p < 0.001$). B) Mutation of the HIV-1^{NL4.3}

capsid (HIV-1^{NL4.3 CA I73V} (MCR) or HIV-1^{NL4.3 CA I73V} (NL4.3)) renders it susceptible to restriction in HeLa-CD4-shREAF cells ($p = 0.03$). C) Fold inhibition of HIV-1^{NL4.3} (VSV-G) with CA mutants I73V and N74D ($p < 0.001$) in the presence of mutant CPSF6-358 compared to vector alone.


Figure 3

Mutant capsids are sensitive to REAF restriction. A) Infection of HeLa-CD4-shREAF cells with HIV-1^{NL4.3 CA N74D} (MCR) renders it susceptible to REAF compared to viral pseudotype with wild type CA (HIV-1^{NL4.3} (MCR)) ($p < 0.001$). B-E) Titres of HIV-1^{NL4.3} (MCR), HIV-1^{NL4.3 CA P38A} (MCR), HIV-1^{NL4.3 CA G89V} (MCR) and HIV-1^{NL4.3 CA G94D} (MCR) following challenge of HeLa-CD4 and HeLa-CD4-shREAF cells. F) Fold change for HIV-1^{NL4.3 CA P38A} (MCR), HIV-1^{NL4.3 CA G89V} (MCR) and HIV-1^{NL4.3 CA G94D} (MCR) on HeLa-CD4-shREAF cells compared to wild type CA (HIV-1^{NL4.3} (MCR)) show they are also susceptible to REAF (all $p < 0.001$).


Figure 4

Viral envelope determines susceptibility to REAF associated restriction. A) Infection of HeLa-CD4-shREAF cells with HIV-1^{NL4.3} (MCN) decreases sensitivity to REAF associated restriction compared to HIV-1^{NL4.3} (MCR) ($p = 0.007$). Comparison of HIV-2^{MCR} and HIV-2^{MCN} Env on pseudotypes carrying B) N74D ($p < 0.001$), C) P38A ($p < 0.001$), D) G89V ($p < 0.001$) and E) G94D ($p = 0.001$) CA show HIV-2^{MCN} Env makes these viruses less sensitive to REAF.


A




B




C




D




E



F




G




2


A



B




C




3


A




B




C




D




E




F




A




B



C



D



E

